

Ichthyofaunal survey of the Riacho Goulart, tributary of the Tietê River (upper Paraná basin)

Marlon F. Pazian^{1*}, Hugo Muelas Moura Leal² and Márcio Lalucce²

1 Universidade Estadual Paulista "Júlio de Mesquita Filho", Instituto de Biociências de Botucatu, Departamento de Morfologia, Distrito de Rubião Junior S/N, CEP 18618-000. Botucatu, SP, Brazil.

2 Faculdade de Filosofia Ciências e Letras de Penápolis, Departamento de Ciências Biológicas, (FAFIPE). Avenida São José 400. CEP 16300-000. Penápolis, SP, Brazil.

* Corresponding author: E-mail: pazian@ibb.unesp.br

ABSTRACT: Fishes assessed in the present study were collected in the Riacho Goulart, a small tributary of Tietê River, upper Paraná basin. In order to sample different habitats, four collection sites were selected. A total of 1,241 individuals belonging to 30 species, 15 families, and six orders were captured. The list of fish species presented herein is important for the knowledge of the biodiversity of low Tietê basin.

INTRODUCTION

South America is the continent with the richest freshwater fish fauna in the world, especially in the Amazon and Paraná-Paraguay river basins. The Amazon basin covers an area of approximately 7,000,000 km², and is estimated to have 2,700 species (Junk *et al.* 2007). The Paraná-Paraguay covers an area of 2,600,000 km² (Latrubesse *et al.* 2005) and has approximately 600 species (Bonetto 1986). The upper Paraná basin covers 900,000 km², and are estimated to have from 130 species (Bonetto 1986) to approximately 310 species (Langeani *et al.* 2007).

On the other hand, in the upper Paraná basin, most rivers, streams and adjacent landscapes are going through extremely diverse changes due to decrease of native vegetation and pollution of waters, which can cause the loss of species, faunal homogenization, and reduction of ichthyic biomass (Casatti 2010). In the northwestern region of São Paulo state, only 3.3% of the natural vegetation remains, arranged in small fragments (mostly up to 10 ha), which characterizes it as one of the most degraded areas of the Brazil (Ferreira and Casatti 2006).

Recent environmental inventories of the headwaters streams of the upper Paraná basin, especially in the state of São Paulo (Casatti *et al.* 2001; Castro and Casatti 1997; Castro *et al.* 2003; 2004; 2005; Langeani *et al.* 2005a, b), have recorded the occurrence of a very diverse fauna, 6-15% of which can be ascribed to undescribed species (Castro *et al.* 2003; 2004; 2005). Similar results were also obtained through the review of ichthyological collections and other independent studies, reinforcing the fact that the surveys in the Upper Paraná are biased (Agostinho and Gomes 2005) and showing the importance of increasing data collection efforts in the area (Langeani *et al.* 2007).

Some fish community surveys have been performed in the lower Tietê basin. Torloni *et al.* (1993) recorded 42 species in the main canal of the Rio Tietê and some of its tributaries. Vidotto and Carvalho (2007) described

the occurrence of 40 species in the rio Santa Bárbara, a tributary of the right margin of the rio Tietê. Ferreira and Casatti (2006) collected 28 species in a tributary of the rio São José dos Dourados. This study provides a list of fish species with the intention of contributing to the knowledge of the biodiversity in upper Paraná basin.

MATERIALS AND METHODS

Study Area

The Riacho Goulart is a small tributary of the Tietê River located approximately 450 meters above sea level (m.a.s.l.). Its headwaters are located in Birigui, São Paulo. Currently, this stream is silted and its margins are mostly occupied by grass, used in cattle farms. The main course of Riacho Goulart has approximately 26 km long. In order to sample diverse number of environments, four collection sites along the Riacho Goulart were selected (Figure 1), **G1** (21°13'00.99" S, 50°21'10.15" W): margins are occupied by grass; **G2** (21°11'36.75" S, 50°20'36.87" W): margins are mostly occupied by grass and few trees; **G3** (21°09'47.56" S, 50°19'31.11" W): margins are mostly occupied by grass and macrophytes; **G4** (21°07'34.55" S, 50°19'23.34" W): margins are mostly occupied by grass and little stones.

Fish sampling

The sampling was done during six months (from December 2004 to May 2005), and two kinds of fishing gear were used: casting-nets (mesh size of 30 mm between adjacent knots, 1,5 m of mouth) and sieve (mesh size of 2 mm, 1 m of length, 0,6 m of depth). In each collection event five times casting-net throws were released (ten meters upstream the place where we used the sieve) and the sieve was used on the bank vegetation in stretches of 30 meters for 60 minutes.

Fishes were fixed with 10% formalin and, after 48 hours, transferred to a 70% ethanol solution. Species

were identified by specialists. Voucher specimens were deposited in the zoological collection of the Laboratório de Zoologia da Faculdade de Filosofia Ciências e Letras Departamento de Ciências Biológicas (FAFIPE), Penápolis, São Paulo and Laboratório de Biologia e Genética de Peixes (LBP), Departamento de Morfologia da Universidade Estadual Paulista (UNESP) Botucatu, São Paulo. The taxonomic classification follows Reis *et al.* (2003) and Langeani *et al.* (2007).

RESULTS AND DISCUSSION

A total of 1,241 individuals belonging to 30 species, distributed in 15 families and six orders, were collected (Table 1). Most species collected in the Riacho Goulart belongs to the orders Characiformes and Siluriformes (76.66%), corroborating the findings reported by Castro *et al.* (2003; 2004), Langeani *et al.* (2007) and Esguícero and Arcifa (2011) for other Neotropical streams.

The present results were coincident with other similar studies for sharing many species reported by them. Thirteen of the 30 species sampled in the Riacho Goulart were also reported by Vidotto and Carvalho (2007) in the Santa Bárbara River. Similarly, 13 of the 28 species reported by Ferreira and Casatti (2006) in the Riacho Água Limpa (São José dos Dourados basin) were also sampled in the Riacho Goulart. These results show that complementary surveys are important for having a comprehensive inventory of a specific area of study.

Finally, in the Riacho Goulart some species sampled were introduced from others basins, including *Satanoperca pappaterra* (Heckel, 1840), originally distributed in the Amazon basin (Kullander 2003), *Roeboides descavadensis* Fowler, 1932 and *Metynnis mola* Eigenmann and Kennedy, 1903 originally distributed in the lower Paraná basin, and introduced after the construction of the Itaipu reservoir (Langeani *et al.* 2007).




FIGURE 1. Hydrographic map of the Riacho Goulart, black circles correspond to sampled sites and geographic coordinates.

TABLE 1. List of the species captured in Riacho Goulart, following the taxonomic classification system by Reis *et al.* (2003) and Langeani *et al.* (2007). G1 corresponds to the first sampled site, G2 corresponds to the second sampled site, G3 corresponds to the third sampled site and G4 corresponds to the fourth sampled site.

TAXON	COLLECTION POINT	VOUCHER
CHARACIFORMES		
Parodontidae		
<i>Parodon nasus</i> Kner, 1859	G4	LBP 9915
Curimatidae		
<i>Cyphocharax modestus</i> (Fernández-Yèpez, 1948)	G1, G2,G4	LBP 11371, 11381
Anostomidae		
<i>Leporinus friderici</i> (Bloch, 1794)	G4	LBP 9916
<i>Leporinus obtusidens</i> (Valenciennes, 1836)	G1, G2,G4	LBP 11373, 9912
Crenuchidae		
<i>Characidium zebra</i> Eigenmann, 1909	G1,G2,G4	LBP 11397
Characidae		
Genera Incertae Sedis in Characidae		
<i>Astyanax altiparanae</i> Garutti and Britski, 2000	G1,G2,G3,G4	LBP 11390, 11383
<i>Hemigrammus</i> cf. <i>marginatus</i> Ellis, 1911	G1	LBP 11376
<i>Moenkhausia intermedia</i> Eigenmann, 1908	G1,G2,G4	LBP 11379
<i>Moenkhausia sanctaefilomenae</i> (Steindachner, 1907)	G1,G2,G3,G4	LBP 11391, 11382
<i>Oligosarcus pinto</i> i Campos, 1945	G4	LBP 11399
<i>Piabina argentea</i> Reinhardt, 1866	G1,G2,G4	LBP 11392, 9909
Characinae		
<i>Roeboides descavadensis</i> Fowler, 1932	G2	LBP 11388
Serrasalminae		
<i>Metynnis mola</i> Eigenmann and Kennedy, 1903	G1,G3	LBP 11374
<i>Serrasalmus marginatus</i> Valenciennes, 1847	G1	LBP 11369
Acestrorhynchidae		
<i>Acestrorhynchus lacustris</i> (Lütken, 1875)	G1	LBP 11368
Erythrinidae		
<i>Hoplerythrinus unitaeniatus</i> (Agassiz, 1829)	G1,G2	LBP 11366
<i>Hoplías malabaricus</i> (Bloch, 1794)	G1,G2,G3,G4	LBP 11386
SILURIFORMES		
Callichthyidae		
<i>Corydoras aeneus</i> (Gill, 1858)	G1,G2,G3,G4	LBP 11377
<i>Hoplosternum littorale</i> (Hancock, 1828)	G3	LBP 11389
Loricariidae		
Hypostominae		
<i>Hypostomus ancistroides</i> (Ihering, 1911)	G1,G2, G3,G4	LBP 11380
<i>Hypostomus</i> cf. <i>paulinus</i> (Ihering, 1905)	G4	LBP 9910, 9914
Heptapteridae		
<i>Pimelodella avanhandavae</i> Eigenmann, 1917	G1,G4	LBP 11398
<i>Rhamdia quelen</i> (Quoy and Gaimard, 1824)	G1,G2,G3	LBP 11372
GYMNOTIFORMES		
Gymnotidae		
<i>Gymnotus</i> cf. <i>carapo</i> Linnaeus, 1758	G1, G2,G3,G4	LBP 9913, 11395
Sternopygidae		
<i>Eigenmannia virescens</i> (Valenciennes, 1847)	G1, G2,G3,G4	LBP 9911, 11394
CYPRINODONTIFORMES		
Poeciliidae		
<i>Phalloceros harpagos</i> Lucinda, 2008	G2,G4	LBP 11393, 11378
SYNBRANCHIFORMES		
Synbranchidae		
<i>Synbranchus marmoratus</i> Bloch, 1795	G1,G2	LBP 11375
PERCIFORMES		
Cichlidae		
<i>Cichlasoma paranaense</i> Kullander, 1983	G1,G2,G3	LBP 11384
<i>Crenicichla britskii</i> Kullander, 1982	G1,G2,G3,G4	LBP 11385, 11396
<i>Satanoperca pappaterra</i> (Heckel, 1840)	G1,G2,G3,G4	LBP 11387

ACKNOWLEDGMENTS: We are grateful to FAFIPE for their support; to Dr Rodrigo Bráz de Castilho Almeida and Dr Artur Antonio Andreata for their collaboration, and to the anonymous reviewers for the suggestions proposed.

LITERATURE CITED

- Agostinho, A.A. and L.C. Gomes. 2005. O manejo da pesca em reservatórios da bacia do Alto rio Paraná: avaliação e perspectivas; p. 23-55 *In* M.G. Nogueira, R. Henry and A. Jorcin (ed.). *Ecologia de reservatórios. Impactos potenciais, ações de manejo e sistemas em cascata*. Volume II. São Carlos: Rima Editora.
- Bonetto, A.A. 1986. The Paraná river system; p. 541-555 *In* B.R. Davies and K.F. Walker (ed.). *The ecology of river systems*. Dordrecht: Dr. W. Junk Publishers.
- Casatti, L. 2010. Changes in the Brazilian Forest Code: potential impacts on the ichthyofauna. *Biota Neotropica* 10(4): 31-34.
- Casatti, L., F. Langeani and R.M.C. Castro. 2001. Peixes de riacho do Parque Estadual Morro do Diabo, bacia do Alto Rio Paraná. *Biota Neotropica* 1(1,2): 1-15.
- Castro, R.M.C. and L. Casatti. 1997. The fish fauna from a small forest stream of the upper Paraná river basin. *Ichthyological Explorartion Freshwaters* 7(4):337-352.
- Castro, R.M.C., L. Casatti, H.F. Santos, K.M. Ferreira, A.C. Ribeiro, R.C. Benine, G.Z.P. Dardis, A.L.A. Melo, T.X. Abreu, F.A. Bockmann, M. Carvalho, F.Z. Gibran and F.C.T. Lima. 2003. Estrutura e composição da ictiofauna de riachos do Rio Paranapanema, sudeste e sul do Brasil. *Biota Neotropica* 3(1): 1-31.
- Castro, R.M.C., L. Casatti, H.F. Santos, A.L.A. Melo, L.S.F. Martins, K.M. Ferreira, F.Z. Gibran, R.C. Benine, M. Carvalho, A.C. Ribeiro, T.X. Abreu, F.A. Bockmann, G.Z.P. Dardis, R. Stopiglia and, F. Langeani. 2004. Estrutura e composição da ictiofauna de riachos da bacia do Rio Grande, no Estado de São Paulo, Sudeste do Brasil. *Biota Neotropica* 4(1): 1-39.
- Castro, R.M.C., L. Casatti, H.F. Santos, R.P. Vari, A.L.A. Melo, L.S.F. Martins, T.X. Abreu, R.C. Benine, F.Z., Gibran, A.C. Ribeiro, F.A. Bockmann, M. Carvalho, G.Z. Pelício, K.M. Ferreira, R. Stopiglia and A. Akama. 2005. Structure and composition of the stream ichthyofauna of four tributary rivers of the upper Rio Paraná basin, Brazil. *Ichthyological Explorartion Freshwaters* 16(3):193-214.
- Esguícero, A.L.H. and M.S. Arcifa. 2011. The fish fauna of the Jacaré-Guaçu river basin, Upper Paraná river basin. *Biota Neotropica* 11(1): 2-11.
- Ferreira, F.C. and L. Casatti. 2006. Integridade biótica de um córrego na bacia do Alto Rio Paraná avaliada por meio da comunidade de peixes. *Biota Neotropica* 6(3): 1-25.
- Junk, W.J., M. G. M. Soares, and P. B. Bailey. 2007. Freshwater fishes of the Amazon River basin: their biodiversity, fisheries, and habitats. *Aquatic Ecosystem Health and Management* 10: 153-173.
- Kullander, S.O. 2003. Family Cichlidae; p. 604-654 *In* R.E. Reis, S.O. Kullander and C.J. Ferraris Jr. (ed.). *Check list of the freshwater fishes of South and Central America*. Porto Alegre: EDIPUCRS.
- Langeani, F., L. Casatti, H.S. Gameiro, A. Bellucco-Do-Carmo and D.C. Rossa-Ferres. 2005a. Riffle and pool fish communities in a large stream of southeastern Brazil. *Neotropical Ichthyology* 3(2): 305-311.
- Langeani, F., Z.M.S. Lucena, J. Pedrini and F.J. Tarelho-Pereira. 2005b. *Bryconamericus turiuba*, a new species from the upper rio Paraná system (Ostariophysi: Characiformes). *Copeia* 2005(2): 386-392.
- Langeani, F., R.M.C. Castro, O.T. Oyakawa, O.A. Shibatta, C. S. Pavanelli and L. Casatti. 2007. Diversidade da ictiofauna do Alto Rio Paraná: composição atual e perspectivas futuras. *Biota Neotropica* 7(3):181-197.
- Latrubesse, E.M., J.C. Stevaux, M.L. Santos and M.L. Assine. 2005. Grandes sistemas fluviais: geologia, geomorfologia e paleohidrologia; p 276-297 *In* C.R.G. Souza, K. Suguio, A.M.S Oliveira and P.E. Oliveira (ed.). *Quaternário no Brasil*. Ribeirão Preto: Editora Holos.
- Reis, R.E., S.O. Kullander and C.J. Ferraris Jr. 2003. *Check list of the freshwater fishes of South and Central America*. Porto Alegre: EDIPUCRS. 742 p.
- Torloni, C.E.C., A.R.A. Corrêa, A.A. Carvalho Jr., J.J. Santos, J.L. Gonçalves, E.J. Gereto, J.A. Cruz, J.A. Moreira, D.C. Silva, E.F. Deus and A.S. Ferreira. 1993. *Produção pesqueira e composição das capturas em reservatórios sob concessão da CESP nos rios Tietê, Paraná e Grande, no período de 1986 a 1991*. São Paulo: Série Pesquisa e Desenvolvimento. 73 p.
- Vidotto, A.P. and E.D. Carvalho. 2007. Composition and structure of fish community in a stretch of the Santa Bárbara river influenced by Nova Avanhandava Reservoir (low Tietê river, São Paulo State, Brazil). *Acta Limnologica Brasiliensia* 19(2): 233-245.

RECEIVED: June 2011

LAST REVISED: September 2011

ACCEPTED: September 2011

PUBLISHED ONLINE: October 2011

EDITORIAL RESPONSIBILITY: Javier A. Maldonado O.